



## ABSTRACT

## Correlation of calorie intake with 24-hour urinary urea nitrogen level changes in critically ill patients at RSUI Depok, Indonesia

Githa Putri Puspita Sari<sup>1</sup>, Fiastuti Witjaksono<sup>1</sup>, Anna Maurina Singal<sup>2</sup>

1. Department of Nutrition, Faculty of Medicine, Universitas Indonesia – Dr. Cipto Mangunkusumo Hospital, Jakarta, Indonesia

2. Department of Nutrition, Faculty of Medicine, Universitas Indonesia – RSUI Depok, Indonesia

Received: 18 October 2024

Accepted: 18 October 2024

Published: 18 October 2024

Link to DOI

[10.25220/WNJ.V08.S1.0041](https://doi.org/10.25220/WNJ.V08.S1.0041)

**Citation:** Sari G P P, Witjaksono F, and Singal A M. Correlation of calorie intake with 24-hour urinary urea nitrogen level changes in critically ill patients at RSUI Depok, Indonesia, World Nutrition Journal. 2024 October 18, 8(S1):42.



**Copyright:** © 2024 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Website

<http://www.worldnutrijournal.org/>

### **Nutri Symposium 2024: Nutrition advancement in healthcare from conception to well-aged perfection: Unveiling nutrition's impact – Oral Presentation**

**Background and objective:** Calorie together with protein intake has been proven to be important factors that play a role in critically ill patients. Hypermetabolism in the early acute phase increases energy expenditure and calorie requirement. Insufficient calorie intake will increase protein degradation for gluconeogenesis, which can be assessed by the level of 24-hour urinary urea nitrogen (UUN). Adequate daily calorie intake is expected to improve the UUN level in patients. This study aims to investigate the correlation of calorie intake with UUN level changes in critically ill patients.

**Methods:** An observational analytical study with a cross-sectional design on critically ill patients at Intensive Care Unit RSUI Depok, aged 18-60 years. A total of 23 participants were included in the study. Calorie intake data were collected on the first 3 days and the level of UUN was assessed on days 3 and 7. Data were analyzed using SPSS to assess the correlation of calorie intake with UUN level changes.

**Results:** Calorie intakes mean from 21 participants on the first 3 days was  $81.57 \pm 27.71\%$ . The mean of UUN levels on day 3 and day 7 were  $8.12 \pm 5.23$  g and  $7.18 \pm 6.56$  g, respectively. There was a decrease in UUN level on the day 7, however, no significant correlation was found in this study between calorie intake and UUN level changes ( $p=0.2$ ).

**Conclusion:** There is no significant correlation between calorie intake on the first 3 days with UUN level changes in critically ill patients.

**Keywords:** calorie intake, urinary urea nitrogen level changes, critically ill patients

#### **Corresponding author:**

Githa Putri Puspita Sari

Department of Nutrition, Faculty of Medicine, Universitas Indonesia – Dr. Cipto Mangunkusumo Hospital, Jakarta, Indonesia

Email: [githa.putri@ui.ac.id](mailto:githa.putri@ui.ac.id)